

# General Mathematics and Computational Science II

## Exercise 19

May 3, 2007

1. Let  $u: [0, 2\pi] \rightarrow \mathbb{R}$  be defined via the Fourier series

$$u(x) = \sum_{k \in \mathbb{Z}} \hat{u}_k e^{ikx} .$$

Show that its derivative  $w(x) = u'(x)$ , assuming it exists, has Fourier coefficients

$$\hat{w}_k = ik \hat{u}_k . \quad (*)$$

2. The previous question suggests a way to differentiate a function  $u: [0, 2\pi] \rightarrow \mathbb{R}$  numerically: First, sample  $u$  on an equidistant grid. Second, compute the Fourier coefficients of its trigonometric interpolant  $v$  as described in class. Third, use (\*) on the Fourier coefficients  $\hat{v}_k$ . Finally, take the inverse DFT to obtain the derivative sampled on the grid.

The problem is that any standard software library DFT assumes that  $j = 0, \dots, N-1$  and  $k = 0, \dots, N-1$  and will store the arrays in precisely this order. Trigonometric interpolation, on the other hand, requires that  $k = -N/2, \dots, N/2-1$ . With which factor, then, must the  $i$ th element, with respect to the order imposed by the software, of the vector  $\hat{v}$  be multiplied so that differentiation is computed correctly?

3. Solve the difference equation

$$\begin{aligned} t_n &= 2t_{n-1} + 2^{n+1} , \\ t_1 &= 0 . \end{aligned}$$