

Partial Differential Equations

Homework 3

due February 28, 2007

1. Prove that the Taylor series with integral remainder for a function $u \in C^N(\mathbb{R}^n)$ is

$$u(x) = \sum_{|\alpha| \leq N-1} \frac{D^\alpha u(x_0)}{\alpha!} (x - x_0)^\alpha + R_N,$$

where

$$R_N = N \sum_{|\alpha|=N} \frac{(x - x_0)^\alpha}{\alpha!} \int_0^1 (1-t)^{N-1} D^\alpha u(x_0 + t(x - x_0)) dt.$$

Hint: Apply the one-dimensional Taylor formula with integral remainder to the function $f(s) = u(x_0 + s(x - x_0))$.

2. In class we have used Liouville's theorem to show that any bounded solution of the Poisson equation

$$-\Delta u = f$$

for $f \in C_c^2(\mathbb{R}^n)$, $n \geq 3$, is given by the solution formula

$$u(x) = \int_{\mathbb{R}^n} \Phi(x - y) f(y) dy$$

up to an arbitrary additive constant. (Evans, p. 30, Theorem 8.)

This statement does not hold in dimension $n = 2$ since solutions are generically unbounded. Use Liouville's theorem to conjecture and prove the corresponding theorem for $n = 2$.

3. Evans, p. 86 problem 5
4. Evans, p. 86 problem 6